



**II Semester M.Sc. Degree Examination, July 2017**  
**(CBCS)**  
**MATHEMATICS**  
**M202T : Complex Analysis**

Time : 3 Hours

Max. Marks : 70

- Instructions :** 1) Answer **any five** questions.  
2) All questions carry **equal** marks.

1. a) Define harmonic functions and evaluate  $\int \frac{z-3}{z^2+2z+5} dz$  where  $C : |z+1-i| = 2$ .

b) Define Bilinear transformation. Find the bilinear transformation which maps the points  $z = 1, i, -1$ , into  $w = i, 0, -i$ .

c) State and prove Morera's theorem. Also evaluate

$$\int_C \frac{e^{2z}}{z^3-1} dz \quad C : |z| = 3. \quad (3+4+7)$$

2. a) State and prove Cauchy's theorem for a rectangle.

b) Let  $f(z)$  be analytic in a region  $G$  with zeros  $a_1, a_2, \dots, a_m$  repeated according to multiplicity. If  $\gamma$  is a simple closed curve in  $G$  which does not pass

through any  $a_k$ , then, prove that  $\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \sum_{k=1}^m n(r : a_k)$ .

c) State and prove Liouville's theorem. (6+5+3)

3. a) Find the radius of convergence of :

i)  $\sum_{n=1}^{\infty} \frac{n!}{n^n} z^n$       ii)  $\sum_{n=0}^{\infty} \frac{z^n}{n!}$ .

b) Define radius of convergence of a power series. If  $R$  is the radius of convergence of  $\sum a_n z^n$ , then prove the following :

- i) The power series converges for  $|z| < R$  and diverges for  $|z| \geq R$ .  
ii) When  $R = 0$ , the power series diverges for all  $z \neq 0 \in \mathbb{C}$  and when  $R = \infty$ , it converges for all  $z \in \mathbb{C}$ .

c) Find the Laurent's series expansion of  $f(z) = \frac{1}{z^2(z-i)}$  in

- i)  $0 < |z| < 1$       ii)  $0 < |z-i| < 1$       iii)  $|z-i| > 1$ .

BMCW (4+6+4)



4. a) State and prove Taylor's theorem.
- b) Let  $z = a$  be an isolated essential singularity of an analytic function  $f(z)$  and  $k = \{ z - a \mid |z - a| < r\}$  be a neighbourhood of 'a'. For a given  $\varepsilon > 0$  and any complex number  $\xi$ , prove that there exists a point  $z$  with  $0 < |z - a| < r$  such that  $|f(z) - \xi| < \varepsilon$ .
- c) Prove that an analytic function comes arbitrarily close to any complex number in the neighbourhood of an essential singularity. (5+6+3)
5. a) Define residue. If  $f(z)$  has a pole of order  $m$  then prove that

$$\text{Res } f(a) = \frac{1}{(m-1)!} \lim_{z \rightarrow a} \frac{d^{m-1}}{dz^{m-1}} \{(z-a)^m f(z)\}$$

- b) Evaluate the following :

$$\text{i) } \int_{-\infty}^{\infty} \frac{x^2}{x^4 + 1} dx \quad \text{ii) } \int_{-\infty}^{\infty} \frac{\sin x}{(x-1)(x^2+1)} dx. \quad \text{(4+5+5)}$$

6. a) Outline the argument principle, and explain why it is called by that name.
- b) Suppose  $f(z)$  and  $g(z)$  are analytic inside and on a closed curve  $V$  and  $|f(z)| > |g(z)| \forall z$ , then show that number of zeros of  $f(z) + g(z)$  equals zeros of  $f(z)$ .
- c) Show that all roots of  $p(z) = z^8 + 4z^3 + 10$  lie between  $1 \leq |z| \leq 2$ . (4+5+5)

7. a) State and prove Phragmen Lindel of theorem.
- b) State and prove Riemann mapping theorem.
- c) Using the result of the Weier-Strass factorization theorem, construct an entire function having zero's at 1, 2, 3. (6+5+3)

8. a) Let  $f(z)$  be analytic in the region  $|z| < \rho$  and let  $z = re^{i\theta}$  be any point of this region. Then prove that

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{(R^2 - r^2) f(Re^{i\theta}) d\phi}{R^2 - 2Rr \cos(\theta + \phi) + r^2}.$$

- b) Derive the Jensen's formula with standard notations. (8+6)